The insulin to carbohydrate ratio should be in the student's school management plan. The ratio is the amount of carbohydrates that will require 1 unit of insulin. For this example, we will use 1 (unit of insulin) : 10 (grams of carbohydrates).

FOOD DOSE EXAMPLE: If the student is having 60 grams of carbohydrates for lunch, then: 60 ÷ 10 = 6 units of rapid-acting insulin for the food dose.

CORRECTION DOSE EXAMPLE: If the student's blood glucose level is 250, the target level is 150 and the correction factor is 100:

1. 250 – 150 = 100

2. 100 ÷ 100 = 1 unit of rapid-acting insulin for the correction dose

TOTAL MEALTIME INSULIN DOSE EXAMPLE: If the student's food dose is 6 and the correction dose is 1:

WITH SYRINGE

PREPPING INSULIN

INJECTING INSULIN



1. **WASH** hands and put on gloves.
2. **PULL**
   out of pen, push it in until it stops.
3. **TAKE**

*This is not a product endorsement from Beaumont Children's Hospital.

CHECK FOR URINE KETONES

IF LEVEL HIGHER THAN 300

Check student's urine for ketones. Use a Keto-Diastix* strip for Ketones testing.

1. **WASH** hands and put on gloves.
2. **COLLECT** fresh urine in a dry, clean cup.
3. **DIP** the test area of the strip in the urine and remove immediately.
4. **WAIT** 15 seconds.

IF KETONES ARE "NEGATIVE" OR "TRACE"

1. **GIVE** at least 8 oz. of water per hour.
2. **RECHECK** ketones at each urination until negative.
3. **CALL** student's parent if ketones are present.

IF KETONES ARE "MODERATE" OR "LARGE"

1. **CONTACT** student's parent.
2. **ENCOURAGE** student to drink water until parent is reached.
3. **STOP** student from any exercise.
4. **CALL 911** for medical assistance (if a parent can't be reached) if student is experiencing abdominal pain, nausea/vomiting or is lethargic.



*This is not a product endorsement from Beaumont Children's Hospital.

SIGNS & SYMPTOMS

Blood glucose level is less than 65. Or if less than 80 with any of the following symptoms:

- **DIZZINESS**
- **NERVOUSNESS**
- **PERSONALITY CHANGE/ IRRATIONAL BEHAVIOR**
- **BLURRY VISION**
- **SHAKINESS**
- **NAUSEA**
- **CRYING**
- **SLUGGISHNESS**
- **SWEATING**
- **POOR COORDINATION**
- **HUNGER**
- **LIGHTHEADEDNESS**
- **IRRITABILITY**
- **DROWSINESS**
- **ERRATIC RESPONSE TO QUESTIONS**
- **INABILITY TO CONCENTRATE**

HYPOGLYCEMIA | Low Blood Sugar

COMMON CAUSES

Too much insulin
Missed or delayed food
Too much or too intense exercise
Unscheduled exercise

SYMPTOMS

MILD

Hunger
Dizziness
Shakiness
Sweating
Lack of concentration
Poor coordination
Personality/behavior change

SEVERE

Loss of consciousness
Seizure
Inability to swallow

ACTION PLAN

BLOOD GLUCOSE <65
OR 65-80 WITH SYMPTOMS

- Provide 15g of carbohydrates, 4 oz juice or 4 glucose tablets
- Wait 15 minutes
- Recheck blood glucose
- Repeat treatment if blood glucose is <80
- If >2 hours before a meal, give a 15g snack of carbohydrates and protein

SEVERE

- Call 911
- DO NOT give anything by mouth
- Contact trained medical personnel
- Roll child on his/her side
- Administer Glucagon as prescribed
- Stay with child
- Contact parents

HYPERGLYCEMIA | High Blood Sugar

COMMON CAUSES

Too little insulin
Too much food
Decreased activity
Illness/infection or stress

SYMPTOMS

MILD

Increased hunger/thirst
Frequent urination
Fatigue/sleepiness
Blurred vision
Stomach pains
Lack of concentration

SEVERE

Nausea/vomiting
Moderate or large ketones
Sweet, fruity breath
Labored breathing
Confused
Unconscious

ACTION PLAN

CHECK URINE KETONES
IF BS >300 OR SYMPTOMS OF SEVERE HYPERGLYCEMIA

NEGATIVE OR TRACE KETONES

- Give extra water
- Allow use of bathroom as needed
- Inform parents of frequent high readings

SMALL KETONES

- Give at least 8 oz. water every hour
- Recheck ketones at next urination
- Child cannot exercise if ketones present
- Call parent

MODERATE TO LARGE KETONES

- Call parent
- Encourage drinking water until